

ON $W_{\infty/2}$ STRINGS**CHRISTOPHER HULL**
and
ZOHORA KHATUN

Department of Physics, Queen Mary and Westfield College,
Mile End Road, London E1 4NS, England.

ABSTRACT

The string theory defined by gauging $\mathcal{W}_{\infty/2}$, the sub-algebra of $\mathcal{W}_{\infty}$ generated by the currents of even spin, is discussed. The critical value of the central charge is calculated using zeta-function techniques and shown to be $c = 1$. A critical string theory is constructed by coupling a free boson to $\mathcal{W}_{\infty/2}$ -gravity and the physical states are analysed.

Any matter system with a $\mathcal{W}$ -algebra symmetry can be coupled, at least classically, to the corresponding $\mathcal{W}$ -gravity to obtain a theory with local $\mathcal{W}$ -symmetry (a higher-spin generalisation of diffeomorphism symmetry). One can then attempt to quantize the theory, and a consistent critical $\mathcal{W}$ -string theory emerges provided that the matter system is chosen so that all anomalies are cancelled by ghost contributions (for a review, see e.g. [1]). This is equivalent to the existence of a nilpotent BRST charge, and requires in particular that the sum of the matter and ghost central charges should be zero. We will be concerned here with algebras such as $\mathcal{W}_\infty$ or $\mathcal{W}_{1+\infty}$ that are generated by an infinite number of currents, and in particular in the algebra $\mathcal{W}_{\infty/2}$, which is the sub-algebra of $\mathcal{W}_\infty$ generated by the currents of even spin. For such infinitely-generated $\mathcal{W}$ -algebras, there are an infinite number of ghosts and the ghost contribution to the central charge is the sum of a divergent series. One approach to this problem is to regulate the series using the zeta-function method, giving a finite critical central charge. For $\mathcal{W}_\infty$ this gives a critical value of the matter central charge given by $c = -2$, and critical $\mathcal{W}_\infty$ -string theories have been investigated in [6]. Remarkably, for the $\mathcal{W}_{\infty/2}$ sub-algebra, the critical central charge is $c = 1$, so that a critical $\mathcal{W}_{\infty/2}$ -string can be constructed by coupling a free boson to $\mathcal{W}_{\infty/2}$ -gravity, with a linearly-realised $\mathcal{W}$ -symmetry, so that there are no matter-dependent anomalies. This is particularly interesting because of recent developments in the study of $c = 1$ matter coupled to ordinary gravity, a system which has been shown to possess a $\mathcal{W}_\infty$ symmetry. It is the purpose of this paper to investigate the critical $\mathcal{W}_{\infty/2}$ -string theory.

The w_∞ algebra [2] is generated by an infinite set of currents w^s , $s = 2, 3, \dots$, of spin s . The modes w_m^s ($m \in \mathbb{Z}$) satisfy the w_∞ algebra

$$[w_m^s, w_n^r] = [(r+1)m - (s+1)n]w_{m+n}^{s+r} \quad (1)$$

which is a Lie algebra without central extension. The $\mathcal{W}_\infty$ algebra [3] is a deformation of this which includes the Virasoro algebra with central charge c , given by

$$[\mathcal{W}_n^r, \mathcal{W}_m^s] = \sum_{q \geq 0} g_{2q}^{rs}(m, n) \mathcal{W}_{m+n}^{r+s-2q} + c_r(m) \delta^{rs} \delta_{m+n,0} \quad (2)$$

The structure constants $g_{2q}^{rs}(m, n)$ and the central terms $c_r(m)$ are completely fixed by the Jacobi identities and take the following form

$$c_r(m) = m(m^2 - 1)(m^2 - 4) \dots (m^2 - (r+1)^2) c_r \quad (3)$$

where the central charges c_r are given by

$$c_r = \frac{2^{2r-3} r! (r+2)!}{(2r+1)!! (2r+3)!!} c \quad (4)$$

The structure constants are given by

$$g_q^{rs}(m, n) = \frac{1}{2(q+1)!} \phi_q^{rs} N_q^{rs}(m, n) \quad (5)$$

where N_q^{rs} is defined by

$$N_q^{rs}(m, n) = \sum_{k=0}^{q+1} (-1)^k \binom{q+1}{k} [r+1+m]_{q+1-k} [r+1-m]_k [s+1+n]_k [s+1-n]_{q+1-k} \quad (6)$$

with $[x]_n = \Gamma(x+1)/\Gamma(x+1-n)$. Finally the ϕ_q^{rs} are given by

$$\phi_q^{rs} = \sum_{k \geq 0} \frac{(-\frac{1}{2})_k (\frac{3}{2})_k (-\frac{q}{2} - \frac{1}{2})_k (-\frac{q}{2})_k}{k! (-r - \frac{1}{2})_k (-s - \frac{1}{2})_k (r+s-q+\frac{5}{2})_k} \quad (7)$$

where $(x)^n = \Gamma(x+n)/\Gamma(x)$

Restricting to the subset of currents of even spin, $w^{(2r)}$ or $\mathcal{W}^{(2r)}$, gives a closed subalgebra of w_∞ or $\mathcal{W}_\infty$ denoted $w_{\infty/2}$ or $\mathcal{W}_{\infty/2}$ respectively [3, 4]. Similarly restricting to the subset of currents w^s , $s = 2, 2+M, 2+2M, 2+3M, \dots$ gives a subalgebra of w_∞ denoted by $w_{\infty/M}$ [4]. These do not appear to give a subalgebra of $\mathcal{W}_\infty$, however, and it is not known whether a consistent non-degenerate algebra with non trivial central extension exists for currents with these spins if $M \neq 2$.

Given a classical matter system with $\mathcal{W}$ -algebra symmetry, it is straightforward to couple to $\mathcal{W}$ -gravity by introducing a gauge field of spin s corresponding to each conserved $\mathcal{W}$ -current of spin s [4]. On quantisation, one gauges away each gauge field of spin s and introduces a ghost $c^{(s)}$ of spin $1-s$ and an anti-ghost $b^{(s)}$ of spin s . In general, there will be anomalies of two types [5]: (i) if the $\mathcal{W}$ -currents do not generate a closed quantum algebra, there will be matter-dependent anomalies (ii) there will in general be universal anomalies, corresponding to central terms in the current algebra. In particular, there will be a conformal (or gravitational) anomaly corresponding to the total central charge of the Virasoro algebra, given by

$$\begin{aligned} c_{TOT} &= c_{MATTER} + c_{GHOST} \\ c_{GHOST} &= \sum_{s=2}^{\infty} c_s \\ c_s &= -2(6s^2 - 6s + 1) \end{aligned} \quad (8)$$

This anomaly will be absent only if the matter contribution cancels against the ghost contribution.

The requirement of anomaly cancellation is equivalent to the existence of a nilpotent BRST charge Q . If the $\mathcal{W}$ -algebra is a Lie algebra (as in the case for algebras considered above) then it is straightforward to construct a BRST charge by standard techniques and check its nilpotence. A necessary condition for this is the vanishing of c_{TOT} .

The case of the $\mathcal{W}_\infty$ algebra was considered in [7, 8]. The ghost central charge is

$$\begin{aligned} c_{GHOST} &= \sum_{s=2}^{\infty} c_s \\ &= -26 - 74 - 146 - \dots \end{aligned} \quad (9)$$

which is divergent. However, it was suggested in [7, 8] that this divergent sum can be regularised using a zeta-function techniques. Recall that the zeta-function is

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s} \quad (10)$$

so that formally one can define $\sum_{n=1}^{\infty} n^r$ (with $r \geq 0$) to be $\lim_{s \rightarrow -r} \zeta(s)$, which is a well-defined finite number. Thus

$$c_{GHOST} \rightarrow -2[6\zeta(-2) - 6\zeta(-1) + \zeta(0)] \quad (11)$$

Using

$$\zeta(0) = -\frac{1}{2}, \quad \zeta(-1) = -\frac{1}{12}, \quad \zeta(-2) = 0 \quad (12)$$

one finds the regularised value

$$c_{GHOST} = 2 \quad (13)$$

So the anomalies cancel if the matter sector has central charge $c = -2$.

In [7] arguments were given to support this rather formal procedure. It was conjectured that this condition was sufficient for the existence of a nilpotent BRST charge and this was checked explicitly for a number of higher spin anomalies.

Consider now the application of these ideas to $\mathcal{W}_{\infty/2}$. Then

$$\begin{aligned} c_{GHOST} &= \sum_{r=1}^{\infty} c_{2r} \\ &= -2 \sum_{r=1}^{\infty} [6(2r)^2 - 6(2r) + 1] \\ &= -2\{24\zeta(-2) - 12\zeta(-1) + \zeta(-1)\} \end{aligned} \quad (14)$$

giving the regularised value $c_{GHOST} = -1$.

Similarly one can ask for the critical value of the central charge for a theory of $\mathcal{W}_{\infty/M}$ gravity. Then zeta-function regularisation gives

$$\begin{aligned} c_{GHOST} &= \sum_{r=0}^{\infty} c_{2+rM} \\ &= (4M^2 - 15M + 13) \end{aligned} \quad (15)$$

Thus if a theory of critical $\mathcal{W}_{\infty/M}$ strings is to exist, the matter sector must have central charge

$$c_{MATTER}(M) = -(4M^2 - 15M + 13)$$

Note that $c_{MAT} < 0$ unless $M = 2$, so that unless $M = 2$, one might expect problems with unitarity.

In the case of $\mathcal{W}_\infty$, $M = 1$, $c_{MAT} = -2$, a non-linear realisation of $\mathcal{W}_\infty$ with $c_{MAT} = -2$ in terms of one boson was constructed in [7] and used to construct a $\mathcal{W}_\infty$ -gravity theory that was formally anomaly-free. The fact that the $\mathcal{W}$ -symmetry was non-linear meant that there were matter-dependent anomalies [5] and the cancellation of these at $c = -2$ was rather involved [6]. Unlike a $\mathcal{W}$ -algebra generated by a finite number of currents (e.g. $\mathcal{W}_N$), $\mathcal{W}_\infty$ also has linear realisations in terms of free bosons. Given D free complex bosons $\phi^i, \bar{\phi}^i$ the set of currents

$$\mathcal{W}^s = B(s) \sum_{r=1}^{s-1} (-1)^r A_r^s : \partial_z^r \phi \partial_z^{s-r} \bar{\phi} : \quad (16)$$

where,

$$B(s) = q^{s-2} \frac{2^{s-3} s!}{(2s-3)!!} \quad (17)$$

and

$$A_r^s = \frac{1}{(s-1)} \binom{s-1}{r} \binom{s-1}{s-r} \quad (18)$$

satisfy the $\mathcal{W}_\infty$ algebra with $c = 2D$ [9].

In the case of the N -free boson realisation of the Virasoro algebra with $c = N$, it is possible to add a background charge term $\alpha_i \partial^2 \phi^i$ to the stress tensor so that the central charge can take general values $c = N + 24\alpha^2$. We have checked that it is not possible to modify the linear bosonic realisation of $\mathcal{W}_\infty$ to obtain a realisation with general central charges by adding higher derivative terms that are linear in the bosonic fields to the currents (16).

If one restricts the D bosons ϕ^i to be real in this construction, then the odd-spin currents given by (16) vanish, $\mathcal{W}^{(2r+1)} = 0$, and one is left with a set of even spin currents generating the $\mathcal{W}_{\infty/2}$ algebra with $c_{MAT} = D$.

In particular, for one boson, $D = 1$, and we have a realisation of $\mathcal{W}_{\infty/2}$ with the critical c -value, $c_{MAT} = 1$. This means that with a matter system of one boson we have a $\mathcal{W}$ -gravity theory for which the gravitational or conformal anomalies cancel against ghost contributions and we conjecture that the same is true for all higher spin anomalies as well. If this is so then we have constructed a theory of critical $\mathcal{W}_{\infty/2}$ -strings.

The operator quantization of this free boson theory is straightforward. On the cylindrical world-sheets with time coordinate τ and periodic space-coordinate σ ,

$$\phi = \phi_+(\tau + \sigma) + \phi_-(\tau - \sigma) \quad (19)$$

and

$$\partial \phi_+(\sigma) = \sum a_n e^{in\sigma}$$

where the modes a_n satisfy

$$[a_n, a_m] = n \delta_{n+m} \quad (20)$$

As usual, we choose a Fock space representation with momentum p ground state $|p\rangle$ satisfying

$$a_0|p\rangle = p|p\rangle, \quad a_n|p\rangle = 0 \quad n > 0 \quad (21)$$

and we normal-order with respect to this vacuum. A state $|\Omega\rangle$ is a highest weight state of $\mathcal{W}_{\infty/2}$ with weights specified by an infinite-dimensional vector $(h^2, h^4, h^6 \dots)$ if it satisfies

$$\begin{aligned} \mathcal{W}_n^{2r}|\Omega\rangle &= 0, & n > 0 \\ (\mathcal{W}_0^{2r} - h^{2r})|\Omega\rangle &= 0 \end{aligned} \quad (22)$$

for certain intercepts h^{2r} .

We next turn to the physical state conditions. For any $\mathcal{W}$ -string theory, a general state is a linear combination of states, each of which can be written as a product of a matter-sector state $|\psi\rangle$ (an element of the ϕ Fock-space in our case) and a ghost-state. For any given ghost-structure, the condition that the state represents a BRST cohomology class is that $|\psi\rangle$ should be a highest-weight state of the appropriate $\mathcal{W}$ -algebra with weights fixed by the choice of ghost structure. For example, for the bosonic string with conventional ghost vacuum, physical states are highest-weight representations of the Virasoro algebra with weight (intercept) $h^2 = 1$, while for the $\mathcal{W}_3$ -string with conventional ghost vacuum, the weights (h^2, h^4) of $(L_0, \mathcal{W}_0)$ are $(4, 0)$ [10]. For $\mathcal{W}$ -algebras generated by a finite number of currents, there are a number of other ways of determining the intercept for L_0 (see [11]), all of which give the same answer. For $\mathcal{W}$ -strings generated by an infinite number of currents, such as $\mathcal{W}_\infty$ and $\mathcal{W}_{\infty/2}$, each method of calculating the weights of a physical state gives a divergent sum, and different methods can give different results, as can different ways of regularising a given sum.

Suppose a given $\mathcal{W}$ -algebra is generated by a set of currents whose spins s form a set $\mathcal{S}$ (e.g. for $\mathcal{W}_{\infty/2}$ we have $\mathcal{S} = \{2, 4, 6, \dots\}$). The corresponding $\mathcal{W}$ -string has ghosts $\{b^{(s)}, c^{(s)} | s \in \mathcal{S}\}$ and the conventional ghost vacuum is given in terms of the BRST-invariant vacuum $|0\rangle$ by

$$|\Omega\rangle = \left(\prod_{s \in \mathcal{S}} \prod_{i=1}^{s-1} c_i^{(s)} \right) |0\rangle \quad (23)$$

The physical state conditions for a state $|\psi\rangle \otimes |\Omega\rangle$ with this ghost structure then imply that the matter sector state $|\psi\rangle$ should be a highest weight state of the $\mathcal{W}$ -algebra with spin-two intercept h^2 given by

$$h^2 = \sum_{s \in \mathcal{S}} \sum_{n=1}^{s-1} n = \sum_{s \in \mathcal{S}} \frac{1}{2} s(s-1) \quad (24)$$

For the $\mathcal{W}_N$ string, $\mathcal{S} = \{2, 3, 4, \dots, N\}$ and $h^2 = 1, 4, \dots$ for $N = 2, 3, \dots$. For the $\mathcal{W}_{\infty+1}$ string,

$$h^2 = \sum_{s=1}^{\infty} \frac{1}{2} s(s-1) = \frac{1}{2} (\zeta(-2) - \zeta(-1)) = \frac{1}{24} \quad (25)$$

For the $\mathcal{W}_\infty$ string the intercept is the same sum but starting from $s = 2$ instead of $s = 1$. This can be evaluated as

$$h^2 = \sum_{s=2}^{\infty} \frac{1}{2} s(s-1) = \sum_{s=1}^{\infty} \frac{1}{2} s(s-1) - 0 = \frac{1}{24} \quad (26)$$

or as

$$h^2 = \sum_{s=2}^{\infty} \frac{1}{2} s(s-1) = \sum_{r=1}^{\infty} \frac{1}{2} r(r+1) = \frac{1}{2} (\zeta(-2) + \zeta(-1)) = -\frac{1}{24} \quad (27)$$

Clearly, organising the series in different ways can give different results. For $\mathcal{W}_{\infty/2}$, we obtain

$$h^2 = \sum_{s \text{ even}} \frac{1}{2} s(s-1) = \sum_{r=1}^{\infty} r(2r-1) = 2\zeta(-2) - \zeta(-1) = \frac{1}{12} \quad (28)$$

Although it is reassuring that the calculation of the spin-two intercepts for $\mathcal{W}_{1+\infty}$ and $\mathcal{W}_{\infty/2}$ involved no rearrangement of the series, it is not clear whether these are trustworthy results. Indeed, in [7], the intercepts for $\mathcal{W}_{1+\infty}$ and $\mathcal{W}_\infty$ were calculated by a different method and different values were obtained ($h^2 = 0$ in both cases). Furthermore, choosing different ghost vacua will give different intercepts and it is possible that for these theories there may be physical states with different ghost structures. As there does not, at present, seem to be a reliable way of fixing the weights for infinite rank $\mathcal{W}$ -algebras, we shall proceed by analysing the one-boson highest weight representations of $\mathcal{W}_{\infty/2}$ for general weights and return to the issue of how these weights are fixed in the $\mathcal{W}_{\infty/2}$ -string at the end.

We shall now look for highest weight Fock-space states $|\Omega\rangle$ satisfying (22) which are eigenstates of momentum and level number

$$a_0|\Omega\rangle = p|\Omega\rangle, \quad N|\Omega\rangle = n|\Omega\rangle \quad (29)$$

for some momentum p , and level n where $N = \sum a_n a_{-n}$. The mode expansion of $\mathcal{W}^{2s}$ gives the operators

$$\mathcal{W}_l^{2s} = (-1)^{s-1} \sum_{k=1}^{2s-1} \Delta_k^{2s} \sum_{n=-\infty}^{\infty} n^{2s-k-1} (l-n)^{k-1} : a_n a_{l-n} : \quad (30)$$

where,

$$\Delta_k^{2s} = (-1)^k B(2s) A_k^{2s}$$

and

$$B(2s) = \frac{2s^{2s-3} (2s)!}{(4s-3)!!}$$

$$A_k^{2s} = \frac{1}{(2s-1)} \binom{2s-1}{k} \binom{2s-1}{2s-k}$$

so that

$$\begin{aligned}
\mathcal{W}_0^{2s} &= 2(-1)^s \sum_{k=1}^{2s-1} B(2s) A_k^{2s} \sum_{n=1}^{\infty} a_{-n} a_n n^{2s-2} \\
&= 2(-1)^s \sum_{k=1}^{2s-1} B(2s) A_k^{2s} (a_{-1} a_1 + 4a_{-2} a_2 + 9a_{-3} a_3 + 16a_{-4} a_4 + \dots)
\end{aligned} \tag{31}$$

Consider first highest weight states of level 0, and momentum p . The constraint

$$(\mathcal{W}_0^2 - h^2)|p\rangle = 0 \tag{32}$$

implies that

$$p^2 = -h^2, \tag{33}$$

while the higher-spin constraints are satisfied only if

$$h^4 = h^6 = h^8 = \dots = 0 \tag{34}$$

This is the case because for higher spin cases the operators $\mathcal{W}_0^{2s}$ for ($s > 1$) do not contain any $a_0 a_0$ term. Thus states of level 0 only exist if

$$h^4 = h^6 = h^8 = \dots = 0$$

with h^2 arbitrary.

Next consider states of level 1, which are of the form $|\Omega\rangle = a_{-1}|p\rangle$. The condition

$$\mathcal{W}_0^2|\Omega\rangle = h^2|\Omega\rangle \tag{35}$$

implies

$$-(p^2 + 2) = h^2 \tag{36}$$

while $\mathcal{W}_1^2|\Omega\rangle = 0$ implies

$$p = 0$$

Thus the only level-one highest weight states must have zero momentum and

$$h^2 = -2 \tag{37}$$

Then

$$\mathcal{W}_1^4|\Omega\rangle = 0, \quad (\mathcal{W}_0^4 - h^4)|\Omega\rangle = 0 \tag{38}$$

only if

$$h^4 = 2^5 \tag{39}$$

For higher-spin, the constraints (22) imply

$$\begin{aligned}
h^{2s} &= 2(-1)^{s-1} \sum_{k=1}^{2s-1} (-1)^{k-1} \Delta_k^{2s} \\
&= 2(-1)^s \sum_{k=1}^{2s-1} \frac{2s^{2s-3}(2s)!}{(4s-3)!!} \cdot \frac{1}{(2s-1)!} \cdot \binom{2s-1}{k} \binom{2s-1}{2s-k}
\end{aligned} \tag{40}$$

For low s, this gives the results

$$h^6 = -2^9, h^8 = 2^{13}, h^{10} = -2^{17}, \dots \tag{41}$$

and leads us to conjecture that the general result is

$$h^{2s} = (-1)^s 2^{4s-3} \tag{42}$$

Thus there are no highest weight states at level one unless the weights h^{2s} take certain fixed values, which we believe to be given by (42). If the weights do take these special values, then there is a physical level-one state with zero momentum.

Now let us consider states of level 2, of the form $|\Omega\rangle = \{\alpha a_{-2} + \beta(a_{-1})^2\}|p\rangle$. This will be a highest weight state provided

$$\begin{aligned}
\mathcal{W}_2^{2s}|\Omega\rangle &= 0 \\
\mathcal{W}_1^{2s}|\Omega\rangle &= 0 \\
\mathcal{W}_0^{2s}|\Omega\rangle &= h^{2s}|\Omega\rangle
\end{aligned} \tag{43}$$

When $s = 1$, (43) implies

$$\begin{aligned}
\mathcal{W}_2^2|\Omega\rangle &= 0 \\
\mathcal{W}_1^2|\Omega\rangle &= 0 \\
\mathcal{W}_0^2|\Omega\rangle &= h^2|\Omega\rangle
\end{aligned} \tag{44}$$

which is equivalent to

$$\begin{aligned}
-(a_1 a_1 + 2a_0 a_2 + \dots)(\alpha a_{-2} + \beta(a_{-1})^2) &= 0 \\
-2(a_0 a_1 + a_{-1} a_2 + \dots)(\alpha a_{-2} + \beta(a_{-1})^2) &= 0 \\
-(a_0 a_0 + 2a_{-1} a_1 + a_2 a_2 + \dots)(\alpha a_{-2} + \beta(a_{-1})^2) &= h^2(\alpha a_{-2} + \beta(a_{-1})^2)
\end{aligned} \tag{45}$$

This implies

$$2p\alpha + \beta = 0 \tag{46}$$

$$2\alpha + \beta p = 0 \tag{47}$$

$$(p^2 + 4) = h^2 \tag{48}$$

The first two equations imply that either $\alpha = \beta = 0$ in which case the state is trivial or

$$p^2 = 1 \quad (49)$$

The final equation in (48) then fixes

$$h^2 = 5 \quad (50)$$

Similarly for the $s = 2$ components of (43), we obtain

$$(\beta - 8p\alpha)(\alpha a_{-2} + \beta a_{-1}^2) = 0 \quad (51)$$

$$(\beta p + 22\alpha)(\alpha a_{-2} + \beta a_{-1}^2) = 0 \quad (52)$$

$$-32(8\alpha a_{-2} + \beta a_{-1}^2)|p > = h^4(\alpha a_{-2} + \beta a_{-1}^2) \quad (53)$$

The first two equations imply that if the state is non-trivial, then

$$p^2 = -\frac{11}{4} \quad (54)$$

This is inconsistent with (49) and so we conclude that there are no physical states at level two.

It is straightforward to give a similar analysis at higher levels and we have checked that there are no physical states at levels 3, 4, or 5. In each case, there are more constraints than there are free variables parameterizing the state at that level, so that the system is over-constrained and there are no solutions. The number of constraints grows with level more rapidly than the number of variables, and we conjecture that there are no further physical states at any higher levels.

We now return to the $\mathcal{W}_{\infty/2}$ -string. For a given ghost vacuum, the physical state conditions imply that the matter part of a physical state should be a highest weight state satisfying (22) with weights

$$h^2 = \bar{h}^2, \quad h^4 = \bar{h}^4, \quad h^6 = \bar{h}^6, \dots \quad (55)$$

for some numbers $\bar{h}^{2r}$ which depend on the choice of ghost vacuum. Our results up to level five imply the following. There is a physical state at level zero only if there is a ghost vacuum that gives intercepts

$$\bar{h}^4 = \bar{h}^6 = \bar{h}^8 = \dots = 0 \quad (56)$$

with $\bar{h}^2$ arbitrary. If so, then the physical state corresponds to a particle in one dimension with momentum $p = \sqrt{-\bar{h}^2}$. Note that (28) suggests $\bar{h}^2 = \frac{1}{12}$ for the standard ghost vacuum. At level one, there are no physical states unless the weights take fixed values

$$\bar{h}^2 = -2, \quad \bar{h}^4 = 2^5, \quad \bar{h}^6 = -2^9, \quad \bar{h}^8 = 2^{13}, \quad \bar{h}^{10} = -2^{17}, \dots \quad (57)$$

in which case there is a single zero-momentum state. For levels 2, 3, 4, 5 there can be no physical states at all, and we conjecture that there are no states at higher levels either. If this is correct, then the spectrum of this $\mathcal{W}_{\infty/2}$ -string is very small indeed: it consists of at most a massive

particle, which will occur only if there is a ghost vacuum which gives intercepts (56), and a zero-momentum state, which will occur only if there is a ghost vacuum which gives intercepts (57). However, deciding whether or not these states actually occur requires a better understanding of the new kind of divergencies that are present in theories with an infinite number of fields than we have at present.

References

- [1] C.M. Hull, Lectures on W-Gravity, W-Geometry and W-Strings, QMW preprint QMW-93-2 (1993), to be published in the Proceedings of the Trieste Summer School in High Energy Physics and Cosmology.
- [2] I. Bakas, Phys. Lett. **228B** (1989) 57.
- [3] C.N. Pope, L.J. Romans and X. Shen, Nucl. Phys. **B339** (1990) 191.
- [4] C.M. Hull, Nucl. Phys. **B353** (1991) 707.
- [5] C.M.Hull, Nucl. Phys. **B67** (1991) 731.
- [6] E. Bergshoeff, P.S. Howe, C.N Pope, E. Sezgin, X. Shen and K.S. Stelle, Nucl. Phys. **B363** (1991) 163
- [7] C.N. Pope L.J.Romans and X. Shen, Phys. Lett. **B254** (1991) 401.
- [8] K.Yamagishi, Phys. Lett. **266B** (1991) 370.
- [9] I.Bakas and E.Kiritsis,Nucl. Phys. **B343** (1990) 189.
- [10] J.Thierry-Mieg, Phys. Lett. **B197** (1987) 368
- [11] A. Bilal and J.-L. Gervais, Nucl. Phys. **B326** (1989) 222.